

Guideline on Acquired Temporomandibular Disorders in Infants, Children, and Adolescents

Originating Committee

Clinical Affairs Committee – Temporomandibular Joint Problems in Children Subcommittee

Review Council

Council on Clinical Affairs

Adopted

1990

Revised

1999, 2002, 2006

Purpose

The American Academy of Pediatric Dentistry (AAPD) recognizes that disorders of the temporomandibular joint occasionally occur in infants, children, and adolescents. This guideline is intended to assist the practitioner in the recognition and diagnosis of temporomandibular disorders (TMD) and to identify possible treatment options. It is beyond the scope of this document to recommend the use of specific treatment modalities.

Methods

This guideline was developed following the AAPD's 1989 symposium on TMD in children and adolescents.¹ This revision is based upon a review of current dental and medical literature related to TMD in children and adolescents. A MEDLINE search was conducted using the terms "temporomandibular disorder", "adolescent", and "children".

Background

All comprehensive dental examinations should include a screening evaluation of the temporomandibular joint (TMJ) and surrounding area.²⁻⁴ Diagnosis of TMD is based upon a combination of historical information, clinical examination, and/or craniocervical and TMJ imaging.⁵⁻⁶ The findings are classified as symptoms (those reported by the child or parent) and signs (those identified by the dentist during the examination).² The reported prevalence of TMD in infants, children, and adolescents varies widely in the literature.⁷⁻¹⁰ Prevalence of signs and symptoms increases with age. One study of the primary dentition reported 34% with signs and/or symptoms of TMD.¹¹ An epidemiological study of 440 children aged 7-14 reported 36% with symptoms (15% of those having recurrent headaches and 13% clicking sounds).¹² More than half of the children (64%) claimed pain upon palpation of the TMJ muscles.¹² In a study of 285 17 year-olds, 62% had either signs or symptoms of dysfunction, with fatigue in the jaw, TMJ sounds, and tenderness in the lateral pterygoid muscle as the most common findings.¹³

Controversy surrounds the significance of signs and symptoms in this age group, the value of certain diagnostic procedures, and what constitutes appropriate therapy. It is not clear whether these signs and symptoms constitute normal variation, preclinical features, or manifestations of a disease state.

Temporomandibular disorders have multiple etiological factors.¹⁴ Many studies show a poor correlation between any single etiological factor and resulting signs and symptoms.¹⁴ Research is insufficient to predict reliably which patient will or will not develop TMD. Etiologic factors suggested as contributing to the development of TMD are:

1. Head and neck trauma: Trauma to the chin, a common occurrence in childhood because of a fall, is reported to be a factor in the development of TMD in pediatric patients.^{15,16} Unilateral and bilateral intracapsular or subcondylar fractures are the most common mandibular fractures in children.^{17,18} Closed reduction and prolonged immobilization can result in ankylosis.¹⁹
2. Occlusal factors: There is a relatively low association of occlusal factors and the development of temporomandibular disorders.^{20,21} However, several occlusal features characterize diagnostic groups²²:
 - skeletal anterior open bite;
 - overjet greater than 6 to 7 mm;
 - retrocuspal position (centric relation) to intercuspal position (centric occlusion) slides greater than 4 mm;
 - unilateral lingual cross bite;
 - 5 or more missing posterior teeth.
3. Parafunctional habits: The literature on the association between parafunction and TMD in pediatric patients is contradictory.²³
4. Posture: Craniocervical posture has been associated with occlusion and with dysfunction of the TMJ, including abnormalities of the mandibular fossa, condyle, ramus, and disc.²⁴⁻²⁶
5. Orthodontic treatment: Current literature does not support that the development of TMD is caused by orthodontic treatment.^{27,28}

Certain medical conditions are reported to mimic TMD occasionally. Among them are trigeminal neuralgia, central nervous system lesions, odontogenic pain, sinus pain, otological pain, developmental abnormalities, neoplasias, parotid diseases, vascular diseases, myofascial pain, cervical muscle dysfunction, and Eagle's syndrome. Other common medical conditions (eg, otitis media, allergies, airway congestion, rheumatoid arthritis) can cause symptoms similar to TMD.²⁹

Few studies document success or failure of specific treatment modalities for TMD in infants, children, and adolescents on a long-term basis. These suggest that simple, conservative, and reversible types of therapy are effective in reducing most TMD symptoms in children.^{30,31} Reversible therapies include patient education, physical therapy, behavioral therapy (eg, eliminating chewing gum), medications, and occlusal splints. Irreversible therapies can include occlusal adjustment, mandibular repositioning, and orthodontics.

Recommendations

Every comprehensive dental history and examination should include a TMJ history and assessment. The history should include questions concerning the presence of head and neck pain and mandibular dysfunction, previous orofacial trauma, and history of present illness with an account of current symptoms. In the presence of a positive history and/or signs and symptoms of TMD, the examination should include palpation of masticatory and associated muscles and the TMJs, documentation of joint sounds, occlusal analysis, and assessment of range of mandibular movements including maximum opening, protrusion, and lateral excursions.

Joint imaging is indicated on a selected basis for joint sounds in the absence of other TMD signs and symptoms. For example, the presence of crepitus may indicate degenerative change that is not yet painful.

Therapeutic modalities to prevent TMD in the pediatric population are yet to be supported by controlled studies. For children and adolescents with signs and symptoms of TMD, reversible therapies should be considered. Because of inadequate data regarding their usefulness, irreversible therapies should be avoided.³²

Referral to a medical specialist may be indicated when otitis media, allergies, abnormal posture, airway congestion, rheumatoid arthritis, or other medical conditions are suspected.

References

1. American Academy of Pediatric Dentistry. Treatment of temporomandibular disorders in children: Summary statements and recommendations. *J Am Dent Assoc* 1990;120(3):265-9.
2. McDonald RE, Avery DR, Dean JA. Examination, diagnosis, and treatment planning. In: *Dentistry for the Child and Adolescent*. 8th ed. St. Louis, Mo: Mosby; 2004:1-22.
3. Casamassimo PS, Christensen JR, Fields HW Jr, Ganzberg S. Examination, diagnosis, and treatment planning for general and orthodontic problems. In: Pinkham JR, Casamassimo PS, McTigue DJ, Fields HW Jr, Nowak AJ, eds. *Pediatric Dentistry: Infancy through Adolescence*. 4th ed. St. Louis, Mo: Elsevier Saunders 2005:661-89.
4. American Academy of Pediatric Dentistry. Guideline on recordkeeping. *Pediatr Dent* 2005;27(suppl):176-81.
5. American Academy of Orofacial Pain. Assessment of orofacial pain disorders. In: Okeson J, ed. *Orofacial Pain: Guidelines for Assessment, Diagnosis, and Management*. Carol Stream, Ill: Quintessence Publishing Co Inc. 1996:19-44.
6. Wahlund K, List T, Dworkin SF. Temporomandibular disorders in children and adolescents: Reliability of a questionnaire, clinical examination, and diagnosis. *J Orofac Pain* 1998;12(1):42-51.
7. Alamoudi N, Farsi N, Salako N, Feteih R. Temporomandibular disorders among school children. *J Clin Pediatr Dent* 1998;22(4):323-9.
8. List T, Wahlund K, Wenneberg B, Dworkin SF. TMD in children and adolescents: Prevalence of pain, gender differences, and perceived treatment need. *J Orofac Pain* 1999;13(1):9-20.
9. Stockstill JW, Bowley JF, Dunning D, Spalding P, Stafford K, Erickson L. Prevalence of temporomandibular disorders in children based on physical signs. *J Dent Child* 1998;65(6):459-67, 438.
10. Paesani D, Salas E, Martinez A, Isberg A. Prevalence of temporomandibular joint disk displacement in infants and young children. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1999;87(1):15-9.
11. Bonjardim LR, Baviao MB, Carmagnani FG, Pereira LF, Castelo PM. Signs and symptoms of temporomandibular joint dysfunction in children with primary dentition. *J Clin Pediatr Dent* 2003;28(1):53-8.
12. Nilner M, Lassing SA. Prevalence of functional disturbances and diseases of the stomatognathic system in 7-14 year olds. *Swed Dent* 1981;5(5-6):173-87.
13. Wanman A, Agerberg G. Relationship between signs and symptoms of mandibular dysfunction in adolescents. *Community Dent Oral Epidemiol* 1986;14(4):225-30.
14. Green CS. Etiology of temporomandibular disorders. *Semin Orthod* 1995;1:222-8.
15. Pullinger AG, Seligman DA. Trauma history in diagnostic groups of temporomandibular disorders. *Oral Surg Oral Med Oral Pathol* 1991;71(5):529-34.
16. Greco CM, Rudy TE, Turk DC, Herlich A, Zaki HH. Traumatic onset of temporomandibular disorders: Positive effects of a standardized conservative treatment program. *Clin J Pain* 1997;13(4):337-47.
17. Kaban LB, Mulliken JB, Murray JE. Facial fractures in children: An analysis of 122 fractures in 109 patients. *Plast Reconstr Surg* 1977;59(1):15-20.

18. Posnick JC, Wells M, Pron GE. Pediatric facial fractures: Evolving patterns of treatment. *J Oral Maxillofac Surg* 1993;51(5):836-44; discussion 844-5.
19. Kaban L. Acquired abnormalities of the temporomandibular joint. In: Kaban L, Troulis M, eds. *Pediatric Oral and Maxillofacial Surgery*. Philadelphia, Pa; Saunders 2004:340-76.
20. Pullinger AG, Seligman DA, Gornbein JA. A multiple logistic regression analysis of the risk and relative odds of temporomandibular disorders as a function of common occlusal features. *J Dent Res* 1993;72(6):968-79.
21. McNamara JA Jr, Seligman DA, Okeson JP. Occlusion, orthodontic treatment, and temporomandibular disorders: A review. *J Orofac Pain* 1995;9:73-90.
22. Widmer CG. Evaluation of Temporomandibular Disorders. In: Krause SL, ed. *TMJ Disorders: Management of the Craniomandibular Complex*. Clinics in Physical Therapy. Vol 18. New York, NY. Churchill Livingstone Inc. 1988: 105-9.
23. Widmalm SE, Christiansen RL, Gunn SM. Oral parafunctions as temporomandibular disorder risk factors in children. *Cranio* 1995;13(4):242-6.
24. Sonnesen L, Bakke B, Solow B. Temporomandibular disorder in relation to craniofacial dimensions, head posture and bite force in children selected for orthodontic treatment. *Eur J Orthod* 2001;20(2):179-92.
25. Kondo E, Nakahara R, Ono M, et al. Cervical spine problems in patients with temporomandibular disorder symptoms: An investigation of the orthodontic treatment effects for growing and nongrowing patients. *World J of Orthodontics* 2002;3:295-312.
26. Motoyoshi M, Shimazaki T, Namura S. Biomechanical influences of head posture on occlusion: An experimental study using finite element analysis. *Eur J Orthod* 2002;24(4):319-26.
27. McNamara JA Jr, Turp JC. Orthodontic treatment and temporomandibular disorders: Is there a relationship? Part 1: Clinical studies. *J Orofac Orthop* 1997;58(2):74-89.
28. Hirata RH, Heft MW, Hernandez B, King GJ. Longitudinal study of signs of temporomandibular disorders (TMD) in orthodontically treated and nontreated groups. *Am J Orthod Dentofacial Orthop* 1992;101(1):35-40.
29. Loos PJ, Aaron GA. Standards for management of the pediatric patient with acute pain in the temporomandibular joint or muscles of mastication. *Pediatr Dent* 1989;11(4):331.
30. Bodner L, Miller VJ. Temporomandibular joint dysfunction in children: Evaluation of treatment. *Int J Pediatr Otorhinolaryngol* 1998;44(2):133-7.
31. Skeppar J, Nilner M. Treatment of craniomandibular disorders in children and young adults. *J Orofac Pain* 1993; 7(4):362-9.
32. Koh H, Robinson PG. Occlusal adjustment for treating and preventing temporomandibular joint disorders. *The Cochrane Database of Systematic Reviews* 2003;1:Art. No. CD003812. DOP: 10.1002/146751858.CD003812.